

# On Certain Admissible Embeddings of L-groups

Geo Kam-Fai Tam

Department of Mathematics, University of Toronto

May 11, 2012

## Abstract

Let  $F$  be a local field and  $E/F$  be a separable extension of degree  $n$ . Regard  $T = \text{Res}_{E/F}\mathbb{G}_m$  as an elliptic maximal torus of  $G = \text{GL}_n$ . We can construct an admissible embedding of L-groups  ${}^L T \hookrightarrow {}^L G$  using Langlands-Shelstad  $\chi$ -data. Such embedding gives rise to an induced representation of the Weil group  $W_F$  of  $F$  from a character of  $W_E$ . The relation between induced representations and admissible embeddings provides a different interpretation of the work of Bushnell-Henniart on the essentially tame local Langlands correspondence.

## 1 Introduction

Let  $F$  be a local field and  $W_F$  be the Weil group of  $F$ . Let  $G$  be  $\text{GL}_n$  as a reductive algebraic group over  $F$ . Let  $E$  be a field extension of degree  $n$ . We may regard  $G(F)$  as the automorphism group of the  $F$ -vector space  $E$  by choosing an  $F$ -basis of  $E$ . Write  $E^\times$  as the  $F$ -point of the algebraic torus  $T = \text{Res}_{E/F}\mathbb{G}_m$ . Therefore the multiplicative action of  $T$  on  $E$  gives rise to an  $F$ -embedding  $T \hookrightarrow G$ .

We consider the dual problem as follows. Let  $\hat{G}$  and  $\hat{T}$  be the dual groups of  $G$  and  $T$ . Let  ${}^L G$  and  ${}^L T$  be the corresponding L-groups. By fixing a maximal torus  $\mathcal{T}$  in  $\hat{G}$ , we ask whether there exists an admissible embedding  ${}^L T \hookrightarrow {}^L G$ , an injective morphism of groups that maps  $\hat{T}$  bijectively onto  $\mathcal{T}$  and the  $W_F$ -component of  ${}^L T$  identically to that of  ${}^L G$ . The answer is affirmative and a construction is given by [LS87].

We introduce the idea briefly as follows. The problem can be shown to be equivalent to ask whether the exact sequence  $1 \rightarrow \mathcal{T} \rightarrow {}^L T \rightarrow W_F \rightarrow 1$  splits, and is therefore equivalent to ask whether the cohomology class  $t = t({}^L T) \in H^2(W_F, \mathcal{T})$  defined by  ${}^L T$  is trivial or not. We can construct a splitting for the class  $t$  using a collection of characters  $\{\chi_\lambda\}_\lambda$  called  $\chi$ -data. Here  $\lambda$  runs through  $\mathcal{R}(G, T)$  the root system of the maximal torus  $T$  in  $G$ , and the character  $\chi_\lambda$  is defined on the multiplicative group of a field extension  $E_\lambda$  over some Galois conjugate of  $E$ . We shall go over the properties of  $\chi$ -data and construct the corresponding admissible embedding in Section 3.

The first main result of this article is a relation between admissible embedding and induced representation of Weil group. The 1-cohomology group  $H^1(W_F, \hat{T}) = \text{Int}(\hat{T}) \backslash \text{Hom}_{W_F}(W_F, \hat{T} \rtimes W_F)$  is isomorphic to  $\text{Hom}(W_E, \mathbb{C}^\times)$  naturally. (This fact is known as Shapiro's Lemma.) Using this fact and Proposition 2.3, the set  $\text{Int}(\mathcal{T}) \backslash \text{AE}({}^L T, {}^L G)$  of  $\text{Int}(\mathcal{T})$ -equivalence classes of admissible embeddings  ${}^L T \rightarrow {}^L G$  is a  $\text{Hom}(W_E, \mathbb{C}^\times)$ -torsor. By specifying suitable isomorphism between these bijective sets, we have the following.

**Theorem 1.1** (Proposition 2.5). *Suppose  $\tilde{\xi} \in \text{Hom}_{W_F}(W_F, \hat{T} \rtimes W_F)$  and  $\chi \in \text{AE}({}^L T, {}^L G)$  correspond to characters  $\xi$  and  $\mu$  in  $\text{Hom}(W_E, \mathbb{C}^\times)$  respectively. The composition  $\chi \circ \tilde{\xi}$  when projected to  $\hat{G} = \text{GL}_n(\mathbb{C})$  is isomorphic to  $\text{Ind}_{W_E}^{W_F}(\xi\mu)$  as representations of  $W_F$ .*

The second main result is a converse of the first one. Suppose now  $\chi \in \text{AE}({}^L T, {}^L G)$  is defined by a collection of  $\chi$ -data  $\{\chi_\lambda\}$ . We can recover the character  $\mu$ , in terms of  $\{\chi_\lambda\}$ , that induces the representation  $\text{Ind}_{W_E}^{W_F}(\xi\mu) \cong \chi \circ \tilde{\xi}$ .

**Theorem 1.2** (Proposition 4.2). *Suppose the admissible embedding  $\chi : {}^L T \rightarrow {}^L G$  is defined by  $\chi$ -data  $\{\chi_\lambda\}$ . The character  $\mu$  in Theorem 1.1 can be taken to be*

$$\mu = \prod_{\text{certain } \lambda} \text{Res}_{E^\times}^{E_\lambda^\times} \chi_\lambda.$$

The proof comes from a comparison between the expression of  $\chi$  given by the recipe in [LS87] and the matrix coefficients of  $\text{Ind}_{W_E}^{W_F}(\xi\mu)$ . What missing in Theorem 1.2 is the set through which  $\lambda$  runs in the product form of  $\mu$ . As suggested by the expression of each factor, we take those roots  $\lambda$  whose corresponding field extensions  $E_\lambda$  contain  $E$ . These roots form a set of representatives of the  $W_F$ -orbits of the root system  $\mathcal{R}(G, T)$ . We will be more specific on these representatives using a double coset expression in Proposition 4.1. We emphasize that such  $\mu$  cannot be arbitrary. For example,  $\mu$  satisfies  $\mu|_{F^\times} = \det \text{Ind}_{W_E}^{W_F} 1_{W_E}$ , as shown in Proposition 4.7.

Finally we give an application on a particular case of the local Langlands correspondence, known as the essentially tame case, established in [BH05, BH10]. Let  $F$  be a non-Archimedean local field of characteristic 0, and  $E$  be a tamely ramified extensions over  $F$  of degree  $n$ . For each admissible character [BH05]  $\xi$  of  $E^\times$ , we introduce a character  ${}_F\mu_\xi$  of  $E^\times$ , called the rectifier of  $\xi$ . Its purpose is to measure the difference between a ‘naive’ version of the local Langlands Correspondence [BH05] and the essentially tame one. In a subsequent article [Tam], we prove that the rectifier admits a factorization of the form in Theorem 1.2 with canonical choices of the characters  $\{\chi_\lambda\}$ . In other words, we can express the essentially tame local Langlands Correspondence by admissible embeddings constructed by  $\chi$ -data.

**Outline of the Article** In Section 2 we study the induced representation of Weil group by admissible embedding of L-groups and prove Theorem 1.1. To construct an admissible embedding in general we need  $\chi$ -data, whose definition and properties are discussed in Section 3. We prove the main result Theorem 1.2 and some related facts in Section 4. Finally in Section 5 we describe providently how admissible embedding is related to the essentially tame local Langlands Correspondence.

**Notations** We fix our notations throughout the article. Let  $H$  be a group and  $K$  be a subgroup of  $H$ . The normalizer of  $K$  in  $H$  is denoted by  $N_H(K)$ . Suppose  $H$  acts on a set  $X$ . For  $h \in H$  and  $x \in X$ , we write  ${}^{h \times} x$ , or simply  ${}^h x$ , for the action of  $h$  on  $x$ . The  $H$ -orbit of  $x \in X$  is denoted by  ${}^H x$ . The collection of all  $H$ -orbits of  $X$  is denoted by  $H \backslash X$ . The set of fixed points is denoted by  $X^H$ . If  $f$  is a map whose domain is  $X$ , we write  ${}^h f(x) = f({}^{h^{-1}} x) = f({}^{(h^{-1})} x)$ . If  $X$  is an abelian group, we denote the set of  $j$ -cocycle of  $H$  with values in  $X$  by  $Z^j(H, X)$ , and the  $j$ -cohomology group by  $H^j(H, X)$ .

Given a field extension  $E/F$  and the corresponding Weil groups  $W_E \subseteq W_F$ , we denote induction  $\text{Ind}_{W_E}^{W_F}$  by  $\text{Ind}_{E/F}$  and restriction  $\text{Res}_{W_E}^{W_F}$  by  $\text{Res}_{E/F}$ . For  $G = \text{GL}_n$  as an  $F$ -group we define

$$\hat{G} = \text{GL}_n(\mathbb{C}) \quad \text{and} \quad {}^L G = \text{GL}_n(\mathbb{C}) \times W_F,$$

namely the dual-group and the  $L$ -group of  $G$ . Given a field extension  $E/F$ , let  $T$  be the  $F$ -torus  $T = \text{Res}_{E/F} \mathbb{G}_m$  with

$$\hat{T} = \text{Ind}_{E/F} \mathbb{C}^\times = (\mathbb{C}^\times)^{[E/F]} \quad \text{and} \quad {}^L T = \hat{T} \rtimes W_F$$

as its dual-group and  $L$ -group. Denote the root system of  $T$  in  $G$  by  $\mathcal{R}(G, T)$  and the corresponding Weyl group by  $\Omega(G, T)$ .

## 2 Induction and Admissible Embedding

Let  $G$  be a connected reductive algebraic group defined and quasi-split over  $F$ . Let  $T$  be a maximal torus of  $G$  also defined over  $F$ .

**Definition 2.1.** An *admissible embedding* from  ${}^L T$  to  ${}^L G$  is a morphism of groups  $\chi : {}^L T \rightarrow {}^L G$  of the form

$$\chi(t \rtimes w) = \iota(t) \bar{\chi}(w) \rtimes w$$

for some injective morphism  $\iota : \hat{T} \rightarrow \hat{G}$  and some map  $\bar{\chi} : W_F \rightarrow \hat{G}$ . □

By expanding  $\chi(s \rtimes v) \chi(t \rtimes w) = \chi((s \rtimes v)(t \rtimes w))$ , we can show that

$$(\text{Int} \bar{\chi}(v))({}^{v \hat{G}} \iota(t)) = \iota({}^{v \hat{T}} t) \quad \text{and} \quad \bar{\chi}(vw) = \bar{\chi}(v) {}^{v \hat{G}} \bar{\chi}(w) \tag{1}$$

for all  $t \in \hat{T}$  and  $v, w \in W_F$ . Hence  $\bar{\chi}$  has image in  $N_{\hat{G}}(\iota(\hat{T}))$ . Conversely if  $\iota$  and  $\bar{\chi}$  satisfy (1), then the map in Definition 2.1 is an admissible embedding. We can rephrase (1) as follows. Let  $N_{\hat{G}}(\iota(\hat{T})) \rtimes W_F$  acts on  $\hat{T}$  by  $x \rtimes w t = \iota^{-1}(\text{Int}(x)(w^{\hat{G}} \iota(t)))$ , then the morphism  $W_F \rightarrow \text{Aut}(\hat{T})$ ,  $w \mapsto w_{\hat{T}}$  factors through

$$W_F \rightarrow N_{\hat{G}}(\iota(\hat{T})) \rtimes W_F, w \mapsto \bar{\chi}(w) \rtimes w.$$

Let  $\mathcal{H}$  be a subgroup of  $\hat{G}$ . Two admissible embeddings  $\chi_1, \chi_2$  are called  $\text{Int}(\mathcal{H})$ -equivalent if there is  $x \in \mathcal{H}$  such that  $\chi_1(t \rtimes w) = (x \rtimes 1)\chi_2(t \rtimes w)(x \rtimes 1)^{-1}$  for all  $t \rtimes w \in {}^L T$ . Using (1) we can show that this condition is equivalent to require an  $x \in \mathcal{H}$  giving  $\bar{\chi}_1(w) = x\bar{\chi}_2(w)^{w^{\hat{G}}}x^{-1}$  for all  $w \in W_F$ .

Let's provide more preliminary setup. By taking a conjugate of  $T$  in  $G$ , which is still denoted by  $T$  for brevity, let  $T$  be contained in a Borel subgroup  $B$  defined over  $F$ . Choose an  $W_F$ -invariant splitting  $(\mathcal{T}, \mathcal{B}, \{\hat{X}_{\alpha}\})$  of  $\hat{G}$  and an isomorphism  $(\hat{T}, \hat{B}) \rightarrow (\mathcal{T}, \mathcal{B})$  whose restriction on  $\hat{T}$  is  $\iota$ . For notation convenience we usually omit  $\iota$  and write  $t = \iota(t) \in \mathcal{T}$  for  $t \in \hat{T}$ , but bear in mind that  $\hat{T}$  and  $\mathcal{T}$  may have inequivalent  $W_F$ -actions.

**Remark 2.2.** We choose splittings on  $G$  and  $\hat{G}$  so that we have a duality on the bases of  $T$  and  $\mathcal{T}$  for explicit computations. For example, we choose a basis of  $\mathcal{T}$  for the construction of the Steinberg section (see Section 2.1 of [LS87]). Our main results would be independent of these choices. For instance, the  $\hat{G}$ -conjugacy class of an admissible embedding is independent of the choices of the Borel subgroup  $B$  containing  $T$  and the splitting  $(\mathcal{T}, \mathcal{B}, \{\hat{X}_{\alpha}\})$  of  $\hat{G}$  (see [LS87] (2.6.1) and (2.6.2)).  $\square$

Langlands and Shelstad constructed a particular form of admissible embedding using  $\chi$ -data (see [LS87] (2.5)). We will give the construction in Section 3. Let's assume such construction for a moment, and denote  $\text{AE}({}^L T, {}^L G, (\hat{T}, \hat{B}) \rightarrow (\mathcal{T}, \mathcal{B}))$ , or just  $\text{AE}({}^L T, {}^L G)$  for simplicity, the set of admissible embeddings  ${}^L T \rightarrow {}^L G$  (associated to the choice of the isomorphism  $(\hat{T}, \hat{B}) \rightarrow (\mathcal{T}, \mathcal{B})$ ). The description of this collection is not difficult.

**Proposition 2.3.** *The set  $\text{AE}({}^L T, {}^L G)$  is a  $Z^1(W_F, \hat{T})$ -torsor, and the set of its  $\text{Int}(\mathcal{T})$ -equivalence classes  $\text{Int}(\mathcal{T}) \backslash \text{AE}({}^L T, {}^L G)$  is an  $H^1(W_F, \hat{T})$ -torsor.*

*Proof.* We fix an embedding  $\chi_0 \in \text{AE}({}^L T, {}^L G)$  and take  $\bar{\chi}_0 : W_F \rightarrow \hat{G}$  as in Definition 2.1. Then for each  $\chi \in \text{AE}({}^L T, {}^L G)$ , the difference  $\bar{\chi}\bar{\chi}_0^{-1}$  is a 1-cocycle of  $W_F$  valued in  $\hat{T}$ , i.e.  $\bar{\chi}\bar{\chi}_0^{-1} \in Z^1(W_F, \hat{T})$ . Indeed for a fixed  $w \in W_F$  both  $\bar{\chi}(w)$  and  $\bar{\chi}_0(w)$  project to the same element in  $\Omega(\hat{G}, \mathcal{T}) = N_{\hat{G}}(\mathcal{T})/\mathcal{T}$ . Using (1) we have that

$$\begin{aligned} \bar{\chi}\bar{\chi}_0^{-1}(vw) &= \bar{\chi}(v)^{v^{\hat{G}}} \bar{\chi}(w)^{v^{\hat{G}}} \bar{\chi}_0(w)^{-1} \bar{\chi}_0(v)^{-1} \\ &= \bar{\chi}(v) \bar{\chi}_0(v)^{-1} v^{\hat{T}} \bar{\chi}(w)^{v^{\hat{T}}} \bar{\chi}_0(w)^{-1} \end{aligned}$$

for all  $v, w \in W_F$ . We can readily verify that the map

$$\text{AE}({}^L T, {}^L G) \rightarrow Z^1(W_F, \hat{T}), \chi \mapsto \bar{\chi}\bar{\chi}_0^{-1}$$

is bijective. From the equality  $t\bar{\chi}(v)^{v^{\hat{G}}}t^{-1}\bar{\chi}_0^{-1}(v) = t\bar{\chi}(v)\bar{\chi}_0^{-1}(v)^{v^{\hat{T}}}t^{-1}$  for all  $t \in \mathcal{T}$ , we know that two embeddings are  $\text{Int}(\mathcal{T})$ -equivalent if and only if the corresponding 1-cocycles differ by a coboundary in  $Z^1(W_F, \hat{T})$ .  $\square$

**Remark 2.4.** For  $G = \text{GL}_n$  we can construct an explicit embedding  ${}^L T \rightarrow {}^L G$ . Choose  $\mathcal{T}$  to be the diagonal subgroup of  $\hat{G}$ . We embed  $\hat{T}$  into  $\hat{G}$  with image  $\mathcal{T}$  and define

$$W_F \rightarrow N_{\hat{G}}(\hat{T}), w \mapsto N(w)$$

the permutation matrix whose assignment is according to the  $W_F$ -action on  $\hat{T} \cong \mathbb{C}^{[E/F]}$ , i.e.  $\text{Int}(N(v))t = v^{\hat{T}}t$  for all  $t \in \hat{T}$ . Clearly the map  ${}^L T \rightarrow {}^L G$ ,  $t \rtimes w \mapsto tN(w) \rtimes w$  defines an admissible embedding.  $\square$

For  $G = \text{GL}_n$  and  $T = \text{Res}_{E/F}\text{G}_m$  we give the main result Proposition 2.5 of this section after the following setup. By Shapiro's Lemma (see the Exercise in [Ser79] VII §5.), we have a special case of Langlands Correspondence for torus

$$\text{Hom}(E^{\times}, \mathbb{C}^{\times}) = H^1(W_F, \hat{T}). \quad (2)$$

The precise correspondence is given as follows. Suppose  $\xi$  is a character of  $E^\times$ . We regard  $\xi$  as a character of  $W_E$  by class field theory [Tat79]. Take a collection of coset representatives  $\{g_1, \dots, g_n\}$  of  $W_E \backslash W_F$ . Define for each  $g_i$  a map  $u_{g_i} : W_F \rightarrow W_E$  given by

$$g_i w = u_{g_i}(w) g(g_i, w) \quad \text{for } g(g_i, w) \in \{g_1, \dots, g_n\}. \quad (3)$$

Then define

$$\tilde{\xi} : W_F \rightarrow \hat{T} \cong \mathbb{C}^n, w \mapsto (\xi(u_{g_1}(w)), \dots, \xi(u_{g_n}(w))).$$

It can be checked that  $\tilde{\xi}$  is a 1-cocycle in  $Z^1(W_F, \hat{T})$ , and different choices of coset representatives give cocycles different from  $\tilde{\xi}$  by a 1-coboundary. Hence the 1-cohomology class of  $\tilde{\xi}$  is defined. By abusing of language, we call  $\tilde{\xi}$  a *Langlands parameter* of  $\xi$ . Moreover, combining Proposition 2.3 and (2) we have

$$\text{Hom}(E^\times, \mathbb{C}^\times) = \text{Int}(\mathcal{T}) \backslash \text{AE}({}^L T, {}^L G). \quad (4)$$

Explicitly, if we have a character  $\mu$  of  $E^\times$ , then define

$$\chi : {}^L T \rightarrow {}^L G, t \rtimes w \mapsto t \begin{pmatrix} \mu(u_{g_1}(w)) & & \\ & \ddots & \\ & & \mu(u_{g_n}(w)) \end{pmatrix} N(w) \times w.$$

Here  $N(w)$  is the permutation matrix as introduced in Remark 2.4. Notice that this bijection is non-canonical. Write  $\text{proj} : {}^L G \rightarrow \hat{G}$ ,  $g \times w \mapsto g$ , which is a morphism of groups because  $G = \text{GL}_n$  splits over  $F$ . Combining the bijections (2) and (4), we have the following result.

**Proposition 2.5.** *Suppose that  $\xi$  and  $\mu$  come from  $\tilde{\xi}$  and  $\chi$  by the bijections (2) and (4). The composition*

$$H^1(W_F, \hat{T}) \times \text{Int}(\mathcal{T}) \backslash \text{AE}({}^L T, {}^L G) \rightarrow \text{Int}(\hat{G}) \backslash \text{Hom}_{W_F}(W_F, {}^L G)$$

*such that  $(\tilde{\xi}, \chi) \mapsto \chi \circ \tilde{\xi}$  gives an isomorphism  $\text{proj} \circ \chi \circ \tilde{\xi} \cong \text{Ind}_{E/F}(\xi\mu)$  as representations of  $W_F$ .*

*Proof.* Choose a suitable basis on the representation space of  $\text{Ind}_{E/F}(\xi\mu)$ . For example, if we realize our induced representation by the subspace of functions

$$\{f : W_F \rightarrow \mathbb{C} \mid f(xg) = \xi\mu(x)f(g) \text{ for all } x \in W_E, g \in W_F\},$$

then we choose those  $f_i$  determined by  $f_i(g_j) = \delta_{ij}$  (Kronecker delta) as basis vectors. The matrix coefficient of  $\text{Ind}_{E/F}(\xi\mu)$  is therefore

$$\begin{pmatrix} \mu\xi(u_{g_1}(w)) & & \\ & \ddots & \\ & & \mu\xi(u_{g_n}(w)) \end{pmatrix} N(w)$$

the same matrix as the image of  $\chi \circ \tilde{\xi}$ . □

**Remark 2.6.** We can recover  $\xi$  from  $\text{Ind}_{E/F}\xi$  as follows. We choose the first  $k$  coset representatives  $g_1 = 1, g_2, \dots, g_k$  to be those in the normalizer  $N_{W_F}(W_E) = \text{Aut}_F(E)$ , and (by choosing suitable basis) consider the matrix coefficient of  $\text{Res}_{E/F} \text{Ind}_{E/F}\xi$ . The first  $k$  diagonal entries are always non-zero and give the characters  $\xi^{g_i}$ . □

### 3 Langlands-Shelstad $\chi$ -data

In this section we recall the construction of admissible embeddings  ${}^L T \rightarrow {}^L G$  given in Chapter 2 of [LS87]. Here  $G$  is a connected reductive algebraic group defined and quasi-split over  $F$  and  $T$  is a maximal torus in  $G$  also defined over  $F$ . Take a maximal torus  $\mathcal{T}$  of  $\hat{G}$  and choose a splitting  $(\mathcal{T}, \mathcal{B}, \{\hat{X}_\alpha\})$  for  $\hat{G}$ . We again emphasize that different choices yield  $\text{Int}(\hat{G})$ -equivalent embeddings. For computational convenience we choose  $\mathcal{T}$  to be the diagonal group and  $\mathcal{B}$  to be the group of upper triangular matrices. The tori  $\hat{T}$  and  $\mathcal{T}$  are isomorphic as groups but with different  $W_F$ -actions.

Recall that the existence of an admissible embedding  $\chi : {}^L T \rightarrow {}^L G$  with restriction  $\hat{T} \rightarrow \mathcal{T}$  is equivalent to the existence of a 1-cocycle  $\bar{\chi} \in Z^1(W_F, N_{\hat{G}}(\mathcal{T}))$  as in (1). We write

$$\omega : W_F \rightarrow \Omega(\hat{G}, \mathcal{T}) \rtimes W_F, w \mapsto \omega(w) = \bar{\omega}(w) \rtimes w$$

such that the action of  $\omega(w)$  on  $t \in \hat{T}$  is the same as  ${}^{w\hat{T}}t$ . We recall in (2.1) of [LS87] the Steinberg section  $n_{\text{St}} : \Omega(\hat{G}, \mathcal{T}) \rightarrow N_{\hat{G}}(\mathcal{T})$  and define

$$n : W_F \rightarrow N_{\hat{G}}(\mathcal{T}) \rtimes W_F, w \mapsto n(w) = \bar{n}(w) \rtimes w := n_{\text{St}}(\bar{\omega}(w)) \rtimes w.$$

The map  $n$  may not be a morphism of groups, yet  $\bar{n}$  satisfies the first equation in (1) in place of  $\bar{\chi}$ . We write

$$t_b(v, w) = n(v)n(w)n(vw)^{-1} = \bar{n}(v)^{v_G} \bar{n}(w) \bar{n}(vw)^{-1}, \quad (5)$$

a 2-cocycle of  $W_F$ , whose values are in  $\{\pm 1\}^n \subseteq \mathcal{T}$  by Lemma 2.1.A of [LS87]. Hence the problem of seeking such  $\bar{\chi}$  is equivalent to looking for a map  $r_b : W_F \rightarrow \hat{T}$  that splits  $t_b^{-1}$ , i.e.

$$r_b(v)^{v\hat{T}} r_b(w) r_b(vw)^{-1} = t_b(v, w)^{-1}. \quad (6)$$

**Remark 3.1.** If we regard a group  $\mathcal{H}$  which appears in the exact sequence  $1 \rightarrow \mathcal{T} \rightarrow \mathcal{H} \rightarrow W_F \rightarrow 1$  as a subgroup of  $N_{\hat{G}}(\mathcal{T}) \rtimes W_F$ , then the problem whether such splitting  $r_b$  exists is equivalent to ask whether the exact sequence above with  $\mathcal{H} \cong {}^L T$  splits or not.  $\square$

The idea in [LS87] to construct such splitting  $r_b$  is to choose a set of characters called  $\chi$ -data, which is defined after the following setup. For each  $\lambda$  in the root system  $\mathcal{R} = \mathcal{R}(G, T)$ , we denote the stabilizers

$$W_{+\lambda} = \{w \in W_F | w\lambda = \lambda\} \quad \text{and} \quad W_{\pm\lambda} = \{w \in W_F | w\lambda = \pm\lambda\},$$

and fixed fields

$$E_{+\lambda} = \bar{F}^{W_{+\lambda}} \quad \text{and} \quad E_{\pm\lambda} = \bar{F}^{W_{\pm\lambda}}.$$

In general,  $E_{+\lambda}$  is a field extension of some conjugate of  $E$ . We call a root  $\lambda$  *symmetric* if  $|E_{+\lambda}/E_{\pm\lambda}| = 2$ , and *asymmetric* otherwise. By definition this symmetry is preserved by the  $W_F$ -action. Let

- (i)  $W_F \backslash \mathcal{R}_{\text{sym}}$  be the set of  $W_F$ -orbits of symmetric roots,
- (ii)  $W_F \backslash \mathcal{R}_{\text{asym}}$  be the set of  $W_F$ -orbits of asymmetric roots, and
- (iii)  $W_F \backslash \mathcal{R}_{\text{asym}/\pm}$  be the set of equivalent classes of asymmetric  $W_F$ -orbits by identifying  ${}^{W_F}\lambda$  and  ${}^{W_F}(-\lambda)$ .

**Definition 3.2.** We define a collection of characters  $\{\chi_\lambda : E_{+\lambda}^\times \rightarrow \mathbb{C}^\times | \lambda \in \mathcal{R}\}$ , called  $\chi$ -data, such that the following conditions hold.

- (i) For each  $\lambda \in \mathcal{R}$ , we have  $\chi_{-\lambda} = \chi_\lambda^{-1}$  and  $\chi_{w\lambda} = \chi_\lambda^{w^{-1}}$  for all  $w \in W_F$ .
- (ii) If  $\lambda$  is symmetric, then  $\chi|_{E_{\pm\lambda}^\times}$  equals the quadratic character  $\delta_{E_{+\lambda}/E_{\pm\lambda}}$  attached to the extension  $E_{+\lambda}/E_{\pm\lambda}$ .

$\square$

If we choose  $\mathcal{R}_0$  to be a subset of  $\mathcal{R}$  consisting of representatives of  $W_F \backslash \mathcal{R}_{\text{sym}}$  and  $W_F \backslash \mathcal{R}_{\text{asym}/\pm}$ , then by condition (i) it is enough to define a collection of  $\chi$ -data on  $\mathcal{R}_0$ . For a chosen  $\chi$ -data  $\{\chi_\lambda\}_{\lambda \in \mathcal{R}_0}$ , following (2.5) of [LS87] we define for each  $\lambda \in \mathcal{R}_0$  a map

$$r_\lambda : W_F \rightarrow \mathcal{T}, \quad w \mapsto \prod_{g_i \in W_{\pm\lambda} \backslash W_F} \chi_\lambda(v_1(u_{g_i}(w)))^{g_i^{-1}\lambda}, \quad (7)$$

where  $u_{g_i}$  is the map (3) for  $W_{\pm\lambda} \backslash W_F$  and  $v_1$  is defined similarly for  $W_{+\lambda} \backslash W_{\pm\lambda}$ . We then define

$$r_g = \prod_{\lambda \in \mathcal{R}_0} r_\lambda. \quad (8)$$

Such construction yields (Lemma 2.5.A of [LS87]) a 2-cocycle

$$t_g(v, w) = r_g(v)^{v\tau} r_g(w) r_g(vw)^{-1} \in Z^2(W_F, \{\pm 1\}^n). \quad (9)$$

In constructing the 2-cocycles (5) and (9) we implicitly used two different notions of gauges (defined just before Lemma 2.1.B of [LS87]) on the set  $\mathcal{R}$ . To relate them we introduce a map (see (2.4) of [LS87])  $s = s_{b/g} : W_F \rightarrow \{\pm 1\}^n$  such that

$$s(v)^{v\tau} s(w) s(vw)^{-1} = t_b(v, w) t_g(v, w)^{-1}. \quad (10)$$

Write  $r_b = s_{b/g} r_g$  and  $\bar{\chi} = r_b \bar{n}$ .

**Proposition 3.3.** *The map  $\chi$  defines an admissible embedding  ${}^L T \rightarrow {}^L G$ .*

*Proof.* It suffices to show that  $\bar{\chi}$  satisfies the two conditions in (1). The first condition is just from the definition of  $n(w)$ , while the second condition is a straightforward calculation using (5), (6), (9), and (10).  $\square$

## 4 The Main Results

Let  $G = \mathrm{GL}_n$  and  $T = \mathrm{Res}_{E/F} \mathbb{G}_m$ , both regarded as algebraic groups over  $F$ . Any root in the root system  $\mathcal{R} = \mathcal{R}(G, T)$  can be expressed as  $\begin{bmatrix} g \\ h \end{bmatrix}$  for some  $g, h \in W_F$ , with

$$\begin{bmatrix} g \\ h \end{bmatrix}(t) = g^{-1} t^{(h^{-1}t)^{-1}}$$

for all  $t \in E^\times$ . The  $W_F$ -action on  $\mathcal{R}$  is given by  ${}^w \begin{bmatrix} g \\ h \end{bmatrix} = \begin{bmatrix} gw^{-1} \\ hw^{-1} \end{bmatrix}$  for  $w \in W_F$ . Notice that such action factors through the one of the Weyl group  $\Omega(G, T)$ . If we choose a collection of coset representatives  $\{g_1 = 1, g_2, \dots, g_n\}$  of  $W_E \backslash W_F$ , then we can write  $\lambda = \begin{bmatrix} g_i \\ g_j \end{bmatrix}$  with  $i \neq j$ . It is clear that each orbit  ${}^{W_F} \lambda$  contains a root of the form  $\begin{bmatrix} 1 \\ g \end{bmatrix}$  for some  $g \in \{g_2, \dots, g_n\}$ .

**Proposition 4.1.** *The set  $W_F \backslash \mathcal{R}$  of  $W_F$ -orbits of the root system  $\mathcal{R}$  is bijective to the collection of non-trivial double cosets in  $W_E \backslash W_F / W_E$ , by*

$$W_F \backslash \mathcal{R} \rightarrow (W_E \backslash W_F / W_E) - \{W_E\}, \quad {}^{W_F} \lambda = {}^{W_F} \begin{bmatrix} 1 \\ g \end{bmatrix} \mapsto W_E g W_E.$$

*Proof.* The set of roots  $\mathcal{R}$  can be identified with the set of off-diagonal elements of  $W_E \backslash W_F \times W_E \backslash W_F$ , with  $W_F$ -action by  ${}^g (W_E g_1, W_E g_2) = (W_E g_1 g^{-1}, W_E g_2 g^{-1})$ . By elementary group theory, we know that the orbits are bijective to the non-trivial double cosets in  $W_E \backslash W_F / W_E$ .  $\square$

We denote  $(W_E \backslash W_F / W_E)'$  the collection of non-trivial double cosets, and  $[g]$  the double coset  $W_E g W_E$ . We call  $g \in W_F$  *symmetric* if  $[g] = [g^{-1}]$  and *asymmetric* otherwise. Clearly such symmetry descends to an analogous property on  $(W_E \backslash W_F / W_E)'$ . By Proposition 4.1 the symmetry of  $(W_E \backslash W_F / W_E)'$  is equivalent to the symmetry of  $W_F \backslash \mathcal{R}$ . Let

- (i)  $(W_E \backslash W_F / W_E)_{sym}$  be the set of symmetric non-trivial double cosets,
- (ii)  $(W_E \backslash W_F / W_E)_{asym}$  be the set of asymmetric non-trivial double cosets, and
- (iii)  $(W_E \backslash W_F / W_E)_{asym/\pm}$  be the set of equivalent classes of  $(W_E \backslash W_F / W_E)_{asym}$  by identifying  $[g]$  with  $[g^{-1}]$ .

Let  $\mathcal{D}$  to be a set of representatives in  $g \in W_E \backslash W_F$  of  $(W_E \backslash W_F / W_E)_{sym}$  and  $(W_E \backslash W_F / W_E)_{asym/\pm}$ . If  $\lambda = \begin{bmatrix} 1 \\ g \end{bmatrix}$ , we write  $\chi_\lambda$  as  $\chi_g$  and  $E_{+\lambda}$  as  $E_g$  which equals  $E_g = g^{-1} E E$ . By condition (i) of Definition 3.2, a collection of  $\chi$ -data  $\{\chi_\lambda\}$  depends only on its sub-collection  $\{\chi_g\}$  for  $g \in \mathcal{D}$ . We also call such sub-collection  $\chi$ -data.

Given  $\chi$ -data  $\{\chi_g\}$  let  $\chi$  be the admissible embedding defined by  $\{\chi_g\}$  as in Proposition 3.3. Let  $\xi$  be a character of  $E^\times$  and  $\tilde{\xi} \in Z^1(W_F, \hat{T})$  be a Langlands parameter of  $\xi$ . (The choice of  $\tilde{\xi}$  is known to be irrelevant.) In Proposition 2.5 we described an induced representation  $\mathrm{Ind}_{E/F} \xi$  as certain embedding of the image of  $\tilde{\xi}$  into  $\mathrm{GL}_n(\mathbb{C})$ . Here we have the reverse.

**Proposition 4.2.** *Given  $\chi$ -data  $\{\chi_g\}$ , define*

$$\mu = \mu_{\{\chi_g\}} = \prod_{[g] \in (W_E \backslash W_F / W_E)'} \text{Res}_{E^\times}^{E_g^\times} \chi_g.$$

*Let  $\chi$  be the admissible embedding defined by  $\{\chi_g\}$ . Then for all character  $\xi$  of  $E^\times$ , the composition*

$$W_F \xrightarrow{\tilde{\chi}} \hat{T} \rtimes W_F \xrightarrow{\chi} \hat{G} \times W_F \xrightarrow{\text{proj}} \text{GL}_n(\mathbb{C})$$

*is isomorphic to  $\text{Ind}_{E/F}(\xi\mu)$  as representations of  $W_F$ .*

**Remark 4.3.** Notice that the product in Proposition 4.2 is uniquely determined by  $\{\chi_\lambda\}$ , i.e. independent of the representative  $g \in \mathcal{D}$ , which is itself a coset representative of  $W_E \backslash W_F$ , of the double coset  $[g]$ . Indeed if  ${}^x \begin{bmatrix} 1 \\ g \end{bmatrix} = \begin{bmatrix} 1 \\ h \end{bmatrix}$  for some  $x \in W_F$ , then  ${}^x E_g = E_h$  and so  $\text{Res}_{E^\times}^{E_g^\times} \chi_h = \text{Res}_{E^\times}^{E_g^\times} \chi_g^{x^{-1}} = \text{Res}_{E^\times}^{E_g^\times} \chi_g$  by condition (i) of Definition 3.2.  $\square$

**Remark 4.4.** Suppose we have fixed a character  $\xi$  of  $E^\times$ . Take a subset  $\{g_1 = 1, g_2, \dots, g_k\}$  of coset representatives of  $W_E \backslash W_F$  in the normalizer  $N_{W_F}(W_E) = \text{Aut}_F(E)$  and write  $\mu_1 = \mu_{\{\chi_g\}}$  as in Proposition 4.2. Then all other characters  $\mu_k$  such that  $\text{Ind}_{E/F}(\xi\mu_k) \cong \text{Ind}_{E/F}(\xi\mu_1)$  are of the form  $\mu_k = \xi^{g_k^{-1}} \mu_1$ . This character  $\mu_k$  also has a factorization in Proposition 4.2 with the same  $\chi$ -data of  $\mu$ , except when  $g = g_k$  the character  $\chi_g$  is changed according to the following.

- (i) If  $g$  is symmetric, then  $\chi_g$  is replaced by  $\xi^{g^{-1}} \chi_g$ .
- (ii) If  $g$  is asymmetric, then  $\chi_g$  is replaced by  $\xi^g \chi_g$  and so  $\chi_{g^{-1}}$  by  $\xi^{-1} \chi_{g^{-1}}$ .

$\square$

*Proof.* (of Proposition 4.2) We first abbreviate  $H = W_F$  and  $K = W_E$ . For each  $\lambda = \begin{bmatrix} 1 \\ g \end{bmatrix}$  we denote  $K_g = K \cap g^{-1}Kg$ , which equals  $W_{+\lambda}$ . If  $[g] \in (K \backslash H/K)_\pm$ , then because  $KgK = Kg^{-1}K$  we can replace  $g$  by an element in  $Kg$  such that  $g^2 \in K$ . Subsequently we have  $g \in W_{\pm\lambda}$  and  $g^2 \in K_g = W_{+\lambda}$ . We denote  $K_{\pm g}$  the group generated by  $K \cap g^{-1}Kg$  and  $g$ , which equals  $W_{\pm\lambda}$ . By condition (i) of Definition 3.2, we rewrite the product in Proposition 4.2 as

$$\prod_{[g] \in (K \backslash H/K)_{\text{asym}/\pm}} (\text{Res}_{E^\times}^{E_g^\times} \chi_g) (\text{Res}_{E^\times}^{E_g^\times} \chi_g^{g^{-1}})^{-1} \prod_{[g] \in (K \backslash H/K)_{\text{sym}}} (\text{Res}_{E^\times}^{E_g^\times} \chi_g). \quad (11)$$

Recall that our dual group  $\mathcal{T}$  is the diagonal subgroup. In order to check  $\chi$  gives rise to a character  $\mu$  as (11) it is enough to consider the first entry of  $r_g$  (see (8) and the discussion in Remark 2.6). From (7) we have

$$r_g(w) = \left( \prod_{[g] \in (K \backslash H/K)_{\text{asym}/\pm}} \prod_{g_i \in K_g \backslash H} \chi_g(u_{g_i}(w))^{\begin{bmatrix} g_i \\ gg_i \end{bmatrix}} \right) \left( \prod_{[g] \in (K \backslash H/K)_{\text{sym}}} \prod_{g_i \in K_{\pm g} \backslash H} \chi_g(v_1 u_{g_i}(w))^{\begin{bmatrix} g_i \\ gg_i \end{bmatrix}} \right).$$

By restricting  $w \in W_E$ , we get the first entry of  $r_g(w)$ , namely

$$r_g(w)_1 = \left( \prod_{g \in (K \backslash H/K)_{\text{asym}/\pm}} \left( \prod_{g_i \in K_g \backslash K} \chi_g(u_{g_i}(w)) \right) \left( \prod_{\substack{g_i \in K_g \backslash H \\ gg_i \in K}} \chi_g(u_{g_i}(w))^{-1} \right) \right) \left( \prod_{g \in (K \backslash H/K)_{\text{sym}}} \left( \prod_{g_i \in K_{\pm g} \backslash K} \chi_g(v_1 u_{g_i}(w)) \right) \left( \prod_{\substack{g_i \in K_{\pm g} \backslash H \\ gg_i \in K}} \chi_g(v_1 u_{g_i}(w))^{-1} \right) \right). \quad (12)$$

We now analysis the products in (12) and match them to those in (11). First, for  $g \in (K \backslash H/K)_{asym/\pm}$ , the first product of (12)

$$\prod_{g_i \in K_g \backslash K} u_{g_i}(w), w \in K$$

is the transfer map  $T_{K_g}^K : K^{\text{ab}} \rightarrow (K_g)^{\text{ab}}$ . By class field theory (see [Tat79]), it corresponds to the inclusion  $E^\times \hookrightarrow E_g^\times$ . Therefore

$$\prod_{g_i \in K_g \backslash K} \chi_g(u_{g_i}(w)) = \text{Res}_{E^\times}^{E_g^\times} \chi_g(w),$$

which is the first factor in (11). Next we consider (the inverse of) the second product of (12)

$$\prod_{\substack{g_i \in K_g \backslash H \\ gg_i \in K}} u_{g_i}(w), w \in K$$

For  $g_i \in K_g \backslash H$  such that  $gg_i \in K$ , we can write  $g_i = g^{-1}x_i$  for some  $x_i$  running through a set in  $K$  of representatives of  $K \cap gKg^{-1} \backslash K$ . If  $u_{x_i}$  is the map (3) for  $K \cap gKg^{-1} \backslash K$  then we have

$$g^{-1}(x_i w) = g^{-1}(u_{x_i}(w) x_{j(x_i, w)}), \quad (13)$$

where  $u_{x_i}(w) \in K \cap gKg^{-1}$ . On the other hand, by regarding  $g^{-1}x_i \in K_g \backslash H$  we have

$$g^{-1}x_i w = u_{g^{-1}x_i}(w) g_{j(g^{-1}x_i, w)} \quad (14)$$

where  $u_{g^{-1}x_i}(w) \in K_g$  and  $g_{j(g^{-1}x_i, w)}$  is of the form  $g^{-1}x_j$  for some  $j$ . By comparing (13) and (14) we have  $g^{-1}u_{x_i}(w)g = u_{g^{-1}x_i}(w)$ . Therefore

$$\prod_i u_{g_i}(w) = g^{-1} \left( \prod_i u_{x_i}(w) \right) g = g^{-1} T_{K \cap gKg^{-1}}^K(w) g$$

and hence

$$\prod_{\substack{g_i \in K_g \backslash H \\ gg_i \in K}} \chi_g(u_{g_i}(w)) = \chi_g^{g^{-1}}(T_{K \cap gKg^{-1}}^K(w)) = (\text{Res}_{E^\times}^{E_g^\times} \chi_g^{g^{-1}})(w)$$

which is (the inverse of) the second factor in (11). Finally, for  $g \in (K \backslash H/K)_{sym}$ , we choose coset representatives  $g_1, \dots, g_k, gg_1, \dots, gg_k$  for  $K_g \backslash H$  such that  $g_1, \dots, g_k$  are those of  $K_{\pm g} \backslash H$ . Moreover we can assume that

$$g_1, \dots, g_h, gg_{h+1}, \dots, gg_{2h} \in K.$$

Hence the third product in (12) is

$$\prod_{\substack{g_i \in K_{\pm g} \backslash H \\ Kg_i = K}} \chi_g(v_1(u_{g_i}(w))) = \prod_{i=1}^h \chi_g(v_1(u_{g_i}(w))). \quad (15)$$

Here  $u_{g_i}$  is the map (3) for  $K_{\pm g} \backslash H$  and so  $v_1 u_{g_i}$  is the one for  $K_g \backslash H$ . For the fourth product in (12), because  $\chi_g^g = \chi_g^{-1}$  (by condition (ii) of Definition 3.2) and  $g(v_1(u_{g_i}(w)))g^{-1} = v_1(u_{gg_i}(w))$ , we have indeed

$$\prod_{g_i \in K_{\pm g} \backslash H, gg_i \in K} \chi_g(v_1 u_{g_i}(w))^{-1} = \prod_{i=h+1}^{2h} \chi_g(v_1(u_{gg_i}(w))). \quad (16)$$

Therefore the product of (15) and (16) is  $\chi_g(T_{K_g}^K(w)) = (\text{Res}_{E^\times}^{E_g^\times} \chi_g)(w)$  which is the last factor of (11).  $\square$



The product  $\mu = \mu_{\{\chi_g\}}$  in Proposition 4.2, as  $\{\chi_g\}$  runs through all  $\chi$ -data, does not produce arbitrary character of  $E^\times$ . Its restriction on  $F^\times$  has a specific form by Proposition 4.7. First recall the following known results. Given a group  $H$  we write  $1_H$  the trivial representation of  $H$ . If  $K$  is a subgroup of  $H$  of finite index, denote  $T_K^H : H^{\text{ab}} \rightarrow K^{\text{ab}}$  the transfer morphism. For any  $g \in H$ , we write  ${}^gK = gKg^{-1}$ .

**Proposition 4.5.** *Let  $\sigma$  and  $\pi$  be finite dimensional representations of  $K$  and  $H$  respectively. We have the following formulae.*

(i) (Mackey's Formula)

$$\text{Res}_K^H \text{Ind}_K^H \sigma \cong \bigoplus_{[g] \in K \backslash H / K} \text{Ind}_{K \cap {}^gK}^K \text{Res}_{K \cap {}^gK}^{{}^gK} ({}^g\sigma).$$

$$(ii) \det \text{Ind}_K^H \sigma \cong (\det \text{Ind}_K^H 1_K)^{\dim \sigma} \otimes (\det \sigma \circ T_K^H).$$

$$(iii) (\det \text{Res}_K^H \pi) \circ T_K^H = (\det \pi)^{|H/K|}.$$

*Proof.* Formulae (i) and (ii) are well-known, for example (i) is proved in [Ser77] 7.3, and (ii) can be found in the Exercise in [Ser79] VII §8. Formula (iii) is direct from (ii) if we take  $\sigma = \text{Res}_K^H \pi$ .  $\square$

In particular, if  $\chi$  is a character of  $K$ , then by (ii) we have

$$\chi \circ T_K^H \cong \left( \det \text{Ind}_K^H \chi \right) \left( \det \text{Ind}_K^H 1_K \right). \quad (17)$$

**Lemma 4.6.** *We have the formula*

$$\det \text{Ind}_K^H 1_K = \prod_{[g] \in (K \backslash H / K)'} \det \text{Ind}_{K_g}^H 1_{K_g}.$$

Notice that  $\det \text{Ind}_{K_g}^H 1_{K_g}$  is independent of the choice of representative  $g$  of the double coset  $[g]$  if we interpret the character as the sign of the canonical  $H$ -action on  $H/K_g$ . For all representatives of  $[g]$ , the corresponding actions are equivalent each other.

*Proof.* (of Lemma 4.6) Applying Mackey's formula on  $\sigma = 1_K$  we obtain

$$\text{Res}_K^H \text{Ind}_K^H 1_K \cong \bigoplus_{[g] \in K \backslash H / K} \text{Ind}_{K_g}^K 1_{K_g}.$$

We take determinant and then transfer morphism  $T_K^H$  on both sides. By (ii) and (iii) of Proposition 4.5 we obtain

$$\left( \det \text{Ind}_K^H 1_K \right)^{|H/K|} = \prod_{[g] \in K \backslash H / K} \left( \det \text{Ind}_{K_g}^H 1_{K_g} \right) \left( \det \text{Ind}_K^H 1_K \right)^{|K/K_g|}.$$

Because the sum of  $|K/K_g|$  for  $[g]$  runs through  $K \backslash H / K$  is  $|H/K|$ , the factor  $\left( \det \text{Ind}_K^H 1_K \right)$  on both sides vanish. What remains gives the desired formula.  $\square$

**Proposition 4.7.** *For all  $\chi$ -data  $\{\chi_g\}$ , if  $\mu$  is the character of  $E^\times$  defined by  $\{\chi_g\}$  as in Proposition 4.2, then  $\mu|_{F^\times} \cong \det \text{Ind}_{E/F} 1_{W_E}$ .*

*Proof.* We first abbreviate  $H = W_F$ ,  $K = W_E$ . For each  $\lambda = \begin{bmatrix} 1 \\ g \end{bmatrix}$  we denote  $K_g = K \cap {}^gK = W_{+\lambda}$  and  $K_{\pm g} = W_{\pm \lambda}$ . The isomorphism in Proposition 4.7 can be rewritten as

$$\prod_{[g] \in (K \backslash H / K)'} \chi_g \circ T_{K_g}^H = \det \text{Ind}_K^H 1_K.$$

By Lemma 4.6 we have to show that

$$\prod_{[g] \in (K \backslash H / K)'} \chi_g \circ T_{K_g}^H = \prod_{[g] \in (K \backslash H / K)'} \det \text{Ind}_{K_g}^H 1_{K_g} \quad (18)$$

By comparing (18) termwise, we claim that

(i) If  $[g] \in (K \backslash H/K)_{asym/\pm}$ , then

$$\left(\chi_g \circ T_{K_g}^H\right) \left(\chi_{g^{-1}} \circ T_{K_{g^{-1}}}^H\right) = \left(\det \text{Ind}_{K_g}^H 1_{K_g}\right) \left(\det \text{Ind}_{K_{g^{-1}}}^H 1_{K_{g^{-1}}}\right) \equiv 1.$$

(ii) If  $[g] \in (K \backslash H/K)_{sym}$ , then  $\chi_g \circ T_{K_g}^H \equiv \det \text{Ind}_{K_g}^H 1_{K_g}$ .

If  $[g] \in (K \backslash H/K)_{asym/\pm}$ , then we have  $K_{g^{-1}} = {}^g K_g$ , which is the stabilizer of the root  $\left[\frac{1}{g^{-1}}\right]$ . Because  $\chi_{g^{-1}} = ({}^g \chi_g)^{-1}$  by condition (i) of Definition 3.2, we have

$$\left(\chi_g \circ T_{K_g}^H\right) \left(\chi_{g^{-1}} \circ T_{K_{g^{-1}}}^H\right) = \left(\chi_g \circ T_{K_g}^H\right) \left({}^g \chi_g^{-1} \circ T_{K_g}^H\right) \equiv 1.$$

On the other hand, since the  $H$ -action on  ${}^H \left[\frac{1}{g}\right] = {}^H \left[\frac{1}{g^{-1}}\right]$  is equivalent to that on  ${}^H \left[\frac{1}{g}\right]$ , we have  $\text{Ind}_{K_g}^H 1_{K_g} \cong \text{Ind}_{K_{g^{-1}}}^H 1_{K_{g^{-1}}}$ . Therefore  $\left(\det \text{Ind}_{K_g}^H 1_{K_g}\right) \left(\det \text{Ind}_{K_{g^{-1}}}^H 1_{K_{g^{-1}}}\right) \equiv 1$ . We have proved the first claim. If  $[g] \in (K \backslash H/K)_{sym}$ , then we have an isomorphism  $\text{Ind}_{K_g}^{K_{\pm g}} 1_{K_g} \cong 1_{K_{\pm g}} \oplus \delta_{K_{\pm g}/K_g}$  as representations of  $K_{\pm g}$ . Here  $\delta_{K_{\pm g}/K_g}$  is the quadratic character of  $K_{\pm g}/K_g$ . We denote this character by  $\delta$ . Hence  $\text{Ind}_{K_g}^H 1_{K_g} \cong \text{Ind}_{K_{\pm g}}^H 1_{K_{\pm g}} \oplus \text{Ind}_{K_{\pm g}}^H \delta$  and

$$\det \text{Ind}_{K_g}^H 1_{K_g} \cong (\det \text{Ind}_{K_{\pm g}}^H 1_{K_{\pm g}})(\det \text{Ind}_{K_{\pm g}}^H \delta) \quad (19)$$

by taking determinant. Now condition (ii) of Definition 3.2, namely  $\chi_g \circ T_{K_g}^{K_{\pm g}} = \delta$ , gives  $\chi_g \circ T_{K_g}^H = \delta \circ T_{K_{\pm g}}^H$ . By (17), this is just the right side of (19). We have proved the second claim and therefore Proposition 4.7.  $\square$

## 5 An application on Local Langlands Correspondence

We recall briefly the essentially tame local Langlands Correspondence, in the sense of [BH05, BH10]. Let  $F$  be a non-Archimedean local field of characteristic 0. Let  $G$  be  $\text{GL}_n$  as a reductive group over  $F$ . Let  $\mathcal{A}_n^{et}(F)$  be the set of the isomorphism classes of irreducible essentially tame supercuspidal representations of  $G(F)$ , and  $\mathcal{G}_n^{et}(F)$  be the set of the equivalence classes of essentially tame  $n$ -dimensional irreducible complex representations of  $W_F$ . The two notions of essential tameness above are defined in [BH05]. These two sets are bijective, whose map

$$\mathcal{L} = \mathcal{L}_n^{et} : \mathcal{G}_n^{et}(F) \rightarrow \mathcal{A}_n^{et}(F)$$

is called the essentially tame local Langlands Correspondence.

We introduce a collection for describing  $\mathcal{L}$  explicitly. Let  $P_n(F)$  be the set of  $W_F$ -equivalence classes  $(E, \xi)$  of admissible characters [BH05]  $\xi$  of  $E^\times$  in which  $E/F$  is a tamely ramified extension of degree  $n$ . By [BH05] we know that  $P_n(F)$  bijectively parameterizes both  $\mathcal{A}_n^{et}(F)$  and  $\mathcal{G}_n^{et}(F)$  simultaneously. Here the bijection  $\sigma : P_n(F) \rightarrow \mathcal{G}_n^{et}(F)$  is simply induction of representations, while the one  $\pi : P_n(F) \rightarrow \mathcal{A}_n^{et}(F)$  is constructed in [BK93] and [BH05]. The ‘naive’ Correspondence  $\pi \circ \sigma^{-1} : \mathcal{G}_n^{et}(F) \rightarrow P_n(F) \rightarrow \mathcal{A}_n^{et}(F)$  does not satisfy certain conditions of the essentially tame local Langlands Correspondence (see Theorem 3.1 of [BH05] or Remark 5.2). In other words, the composition

$$\mu : P_n(F) \xrightarrow{\sigma} \mathcal{G}_n^{et}(F) \xrightarrow{\mathcal{L}} \mathcal{A}_n^{et}(F) \xrightarrow{\pi^{-1}} P_n(F)$$

does not give the identity map on  $P_n(F)$ . In [BH10] it is proved that for each admissible character  $\xi$  of  $E^\times$ , there is a character  ${}_F \mu_\xi$  of  $E^\times$ , called the *rectifier* of  $\xi$ , such that  ${}_F \mu_\xi \xi$  is also admissible and  $\mu(E, \xi) = (E, {}_F \mu_\xi \xi)$ .

In terms of admissible embeddings  ${}^L T \rightarrow {}^L G$ , it means that for each  $\xi$  we have to embed the image of the chosen Langlands parameter of  $\xi$  not by the canonical one defined by the Weyl group action (as in Remark 2.4) but the one twisted by its rectifier  ${}_F \mu_\xi$ . The rectifier  ${}_F \mu_\xi$  is explicitly described in [BH10], and so is the Correspondence  $\mathcal{L}$ . Using this description we prove the following result in [Tam].

**Theorem 5.1.** *For each admissible  $\xi$ , the rectifier  ${}_F\mu_\xi$  has a factorization of the form as in Theorem 1.2, for some canonical choice of  $\chi$ -data.*

The proof requires a substantial amount of new concepts and computations, so it is better to deal with these in a separated article. This Theorem suggests that the rectifier has more properties inherited from that of  $\chi$ -data. Indeed the symmetric structure of  ${}_F\mu_\xi$  form that of  $\chi$ -data  $\{\chi_g\}$  is almost trivial because it is known [Tam] that those  $\chi_g$  are of order at most 2 except for exactly one whose has order at most 4. We will have a closer look on this and also other inherited properties of  ${}_F\mu_\xi$  in [Tam]. The following is a property which can be stated with the knowledge of this article.

**Remark 5.2.** Suppose  $\sigma \in \mathcal{G}_n^{et}(F)$  and  $\pi = \mathcal{L}(\sigma) \in \mathcal{A}_n^{et}(F)$ . Let  $\omega_\pi$  be the central character of  $\pi$ . One of the conditions of Langlands Correspondence, namely  $\omega_\pi = \det \sigma$ , implies that  ${}_F\mu_\xi|_{F^\times} = \det \text{Ind}_{E/F} 1_{W_E}$ . This is a general fact about the restriction of the product of the characters in any  $\chi$ -data as in Proposition 4.7, if we have established Theorem 5.1 beforehand.  $\square$

## 6 Acknowledgements

This article is part of the thesis problem assigned to the author by James Arthur, based on a suggestion that originated with Robert Kottwitz. The author would thank James Arthur for his guidance and support. He would also thank Bin Xu for reading the first draft and for his useful comments. Finally but not least, he would thank Alice Lai for typing this article.

## References

- [BH05] Colin J. Bushnell and Guy Henniart. The essentially tame local Langlands correspondence. I. *J. Amer. Math. Soc.*, 18(3):685–710 (electronic), 2005.
- [BH10] Colin J. Bushnell and Guy Henniart. The essentially tame local Langlands correspondence, III: the general case. *Proc. Lond. Math. Soc. (3)*, 101(2):497–553, 2010.
- [BK93] C.J. Bushnell and P.C. Kutzko. *The admissible dual of  $GL(N)$  via compact open subgroups*. Annals of mathematics studies. Princeton University Press, 1993.
- [LS87] R. P. Langlands and D. Shelstad. On the definition of transfer factors. *Math. Ann.*, 278(1-4):219–271, 1987.
- [Ser77] Jean-Pierre Serre. *Linear representations of finite groups*. Springer-Verlag, New York, 1977. Translated from the second French edition by Leonard L. Scott, Graduate Texts in Mathematics, Vol. 42.
- [Ser79] Jean-Pierre Serre. *Local fields*, volume 67 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1979. Translated from the French by Marvin Jay Greenberg.
- [Tam] Geo Kam-Fai Tam. Admissible embedding of L-groups and essentially tame local Langlands correspondence. (*preprint*).
- [Tat79] J. Tate. Number theoretic background. In *Automorphic forms, representations and L-functions (Proc. Sympos. Pure Math., Oregon State Univ., Corvallis, Ore., 1977), Part 2*, Proc. Sympos. Pure Math., XXXIII, pages 3–26. Amer. Math. Soc., Providence, R.I., 1979.